Providing abortion care using mobile phones: the South African experience


Deborah Constant
Women's Health Research Unit, University of Cape Town,
South Africa
deborah.constant@uct.ac.za

Dr Marijke Alblas
South Africa
malblas@iafrica.com

FIAPAC, Slovenia 3-4 October, 2014



The reach of mhealth (1)...




Source: GSMA mhealth: mhealth feasibility South Africa 2014

The reach of mhealth (2)...

The case for mHealth remains compelling



Al




Source: GSMA mhealth: Understanding the needs and wants of pregnant women and mothers. South Africa July 2014. Consumer report


The reach of mhealth (3)...

South Africa: Shared access reduces the effect of rural/urban divide

Large difference in subscribers between urban and rural areas




Considerable shared access – may continue in rural communities for some time



Source: AMPS, GSMA-MDI Analysis

Restricted - Confidential Information ⊚ GSMA 2013

Phone usage and willingness to pay




Source: GSMA mhealth: Understanding the needs and wants of pregnant women and mothers. South Africa July 2014. Consumer report

Opportunities......



New ideas





m-ASSIST Project (2011-2012)

SMSs coaching through MA


2 weeks, from 1st appt

Self-assessment of abortion completion via mobile

2-5 questions SMS request to do SA sent on Day 11

FP info:

- SMSs
- mobisite: m.ichoosewhen.org.za
- Available on Mxit (cheap+ data light)






Hello! Just a reminder to take the pills you were given, 24 - 48 hours (1-2 days) after u took the pills at the clinic. Put a pill in each cheek and 2 under the tongue, and let them dissolve (break down) there. Take them without water! U can drink 30 min after.

Day 13 SMS

Hi hope you're good. You may still be spotting (a bit of bleeding or brown bits). If you're bleeding like a normal period or more, make sure you tell them this at your clinic appointment

SMS SUPPORT: OUTCOMES AT FOLLOW-UP CLINIC VISIT

Intervention N = 197; Control N = 184



SMS SUPPORT: FEEDBACK


- 99% would recommend the SMS support to a friend undergoing MA
- 98% said the SMSs helped them through the MA process

"I always knew what is going to happen so that kept me going because if it was not for the SMSs I would have come back after 2 days. So they helped me a lot because I didn't even call the clinic. They were my hope."


> "Sometimes the SMSs comforted me. I felt the SMSs understood what I was going through. Felt like a friend



Self-Assessment



SELF-ASSESSMENT USAGE (COMPUTER LOG) AND SA RESULTS VS PROVIDER ASSESSMENT (AT FOLLOW-UP)





M-ASSIST STUDY CONCLUSIONS

Aim: Can mobile phones improve medical abortion delivery by providing:

Support information and reminder
 SMSs during the abortion process

YES

 A self-administered questionnaire to assess completion of the procedure

ONLY PARTIALLY

 Information reinforcing family planning methods by SMS and on a mobisite

YES



LIMITATIONS, BARRIERS AND CHALLENGES





Training needed for SA / registration









New study M-ASSIST+ FU at clinic or over lmage: Validation of SA phone LSUPT vs provider: result LSUPT + need for care Paper checklist I choose when.co **Support** za **SMSs** FΡ decision Enrollment making? at clinic visit 1

ichoose when health info -

STORIES

GET HELP

ABOUT

CONTACT

Welcome!

We hope this info will help you take control...of when and if to have a baby.

You can fall pregnant from having sex just once with no contraception, if it happens during the womans fertile time - and remember, sperm can live in a woman's body for up to 5 days. Take control and prevent an unwanted pregnancy!

Before you read on, note: only condoms prevent sexually transmitted infections (STIs) including HIV. So when using other contraceptive methods AND condoms you have the best protection.







Contraception

Contraception 90 (2014) 226-233

Original research article

Mobile phone messages to provide support to women during the home phase of medical abortion in South Africa: a randomised controlled trial ★★★

Deborah Constant^{a,b,*}, Katherine de Tolly^c, Jane Harries^a, Landon Myer^b

^aWomen's Health Research Unit, School of Public Health and Family Medicine, University of Cape Town, South Africa ^bDivision of Epidemiology & Biostatistics, School of Public Health and Family Medicine, University of Cape Town, South Africa ^cCell-life, Cape Town South Africa

Received 18 October 2013; revised 16 April 2014; accepted 16 April 2014

sept 2014

JMIR MHEALTH AND UHEALTH

de Tolly & Constant

Original Paper

Integrating Mobile Phones into Medical Abortion Provision: Intervention Development, Use, and Lessons Learned From a Randomized Controlled Trial

Katherine Marianne de Tolly¹, BA, Hons, MPhil; Deborah Constant², MPH, MSc

²Women's Health Research Unit, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa



2014 | VOI. 2 | iss.

Assessment of completion of early medical abortion using a text questionnaire on mobile phones compared to a self-administered paper questionnaire among women attending four clinics, Cape Town, South Africa

Deborah Constant, a Katherine de Tolly, b Jane Harries, Landon Myerd



Where to read about it?

¹Cell-Life, Cape Town, South Africa

THANK YOU

- FIAPAC
- Safe Abortion Action Fund
- Department of Reproductive Health and Research,
 World Health Organization
- Women's Health Research Unit, UCT
- Karolinska Institutet Prof K Gemzell Danielsson
- Cell Life, Cape Town